

Visit
FREETESTPAPER.com
for more papers



Website: freetestpaper.com



[Facebook.com/freetestpaper](https://www.facebook.com/freetestpaper)



[Twitter.com/freetestpaper](https://www.twitter.com/freetestpaper)



**JUNYUAN SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2020
SECONDARY FOUR EXPRESS**

CANDIDATE NAME

CLASS

4

E

INDEX NUMBER

CHEMISTRY

6092

PAPER 1

2 September 2020

1 hour

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in.

Write in soft pencil.

Do not use paper clips, glue or correction fluid.

There are forty questions on this paper. Answer all questions. For each question there are four possible answers, A, B, C and D.

Choose the one you consider correct and record your choice in soft pencil on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

A copy of the Periodic Table is printed on page 19.

The use of an approved scientific calculator is expected, where appropriate.

This document consists of 19 printed pages.

1 Which substance would diffuse most quickly?

- A carbon dioxide at 0 °C
- B carbon dioxide at 25 °C
- C neon at 0 °C
- D neon at 25 °C





2 A student tested a solution by adding aqueous sodium hydroxide. A precipitate was not seen because the reagent was added too quickly.

What could not have been present in the solution?

- A Al^{3+} B Ca^{2+} C NH_4^+ D Zn^{2+}


3 Four gas jars each contains one of the gases, ammonia, chlorine and oxygen. A strip of damp blue litmus paper and a strip of damp red litmus paper are placed in each jar.

In which gas jar will both the damp blue litmus paper and the damp red litmus paper change colour?



- 4 The diagram shows the chromatogram obtained by analysis of a single dye.

Three measurements are shown.



How is the R_f value of the dye calculated?

- A $\frac{x}{x+y}$ B $\frac{y}{x+y}$ C $\frac{x}{x+y+z}$ D $\frac{y}{x+y+z}$
- 5 The apparatus shown in Fig. 5.1 can be used to separate pure water from seawater.




Fig. 5.1

Two samples are taken, one at point X and another at point Y.

Which of the following statements about X and Y is incorrect?

- A X and Y can be separated into their components by physical methods.
 B When heated to dryness, X leaves a residue while Y does not.
 C X boils over a range of temperatures, while Y boils at 100 °C.
 D X is a mixture while Y is a compound.

- 6 The following statement describes substance X.

A chemical reaction takes place and heat is liberated when this white solid, X is formed.

Which of the following correctly classifies X and explains why X is classified as such?

	classification	explanation
A	element	When decomposition takes place, an element, X, is produced.
B	compound	Bond forming takes place to produce X.
C	mixture	The reactants and the product X, form a mixture.
D	either an element or a compound	A chemical reaction can produce either an element or a compound.


- 7 The atoms $^{64}_{29}\text{Cu}$ and $^{65}_{30}\text{Zn}$ have the same

- A nucleon number.
- B number of electrons.
- C number of neutrons.
- D proton number.

- 8 Which of the following correctly describes what happens when calcium atoms form calcium ions?


	calcium atoms	ionic equation for the formation
A	gain electrons	$\text{Ca} + 2\text{e}^- \rightarrow \text{Ca}^{2+}$
B	gain electrons	$\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$
C	lose electrons	$\text{Ca} + 2\text{e}^- \rightarrow \text{Ca}^{2+}$
D	lose electrons	$\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$

- 9 Element X has a lattice of positive ions and a 'sea of electrons'.




Which property will X have?

- A It conducts electricity by the movement of ions and electrons.
 B It has a high melting point.
 C It is decomposed by an electric current.
 D It is not malleable.
- 10 When a covalent substance in liquid state boils, its molecules become more widely spaced.
- Which property of the molecules has the most influence on the amount of energy required to boil a covalent substance?
- A the forces of attraction between the molecules
 B the reactivity of the molecules
 C the shape of the molecules
 D the strength of the covalent bonds in the molecules
- 11 The diagrams show the structures of two forms of carbon.



S



T

Which set of data is correct for these two structures?

	conducts electricity	very hard material	can be used as lubricant
A	T	T	S
B	S	T	S
C	S	S	T
D	T	S	T

12 Which statement about ionic compounds is correct?

- A Ionic compounds conduct electricity when solid because they contain charged particles that can move.
- B Ionic compounds consist of a lattice of positive ions and negative ions.
- C Most ionic compounds are solids at room temperature because of the strong attraction between electrons and positive ions.
- D When molten or in aqueous solution, ionic compounds conduct electricity because they contain electrons that can move.

13 Sulfur and selenium (Se) are in the same group of the Periodic Table.

From this, we would expect selenium to form compounds having the formulae

- A SeO , Na_2Se and NaSeO_4 .
- B SeO_2 , Na_2Se and NaSeO_4 .
- C SeO_2 , Na_2Se and Na_2SeO_4 .
- D SeO_3 , NaSe and NaSeO_4 .

14 Students give their own special symbols to five non-metallic elements. All five non-metals are in the same group of the Periodic Table. These non-metals exist as coloured elements.

The special symbols are shown in Fig. 14.1. The order of chemical reactivity of these non-metals is also shown.




Fig. 14.1

A solution containing Cc ions can be displaced by two of the elements in the Group.

Which of the following correctly shows the ionic equation for one such reaction?

- A $\text{Aa}(\text{aq}) + \text{Cc}^-(\text{aq}) \rightarrow \text{Aa}^-(\text{aq}) + \text{Cc}(\text{aq})$
- B $\text{Bb}_2(\text{aq}) + 2\text{Cc}^-(\text{aq}) \rightarrow 2\text{Bb}^-(\text{aq}) + \text{Cc}_2(\text{aq})$
- C $\text{Dd}(\text{aq}) + \text{Cc}^-(\text{aq}) \rightarrow \text{Dd}^-(\text{aq}) + \text{Cc}(\text{aq})$
- D $\text{Ee}_2(\text{aq}) + 2\text{Cc}^-(\text{aq}) \rightarrow 2\text{Ee}^-(\text{aq}) + \text{Cc}_2(\text{aq})$

- 15 The table shows some of the properties of four elements.

Which element is most likely to be a transition metal?

	melting point / °C	density / g/cm ³	electrical conductivity
A	3550	3.5	poor
B	1860	7.2	good
C	660	2.7	good
D	232	7.3	good

- 16 The relative formula masses of four compounds are given.

A student has a 1.0 g sample of each compound.

Which sample contains the highest number of moles of oxygen atoms?


	compound	relative formula mass
A	Al ₂ O ₃	102
B	CuO	80
C	H ₂ SO ₄	98
D	HNO ₃	63

- 17 What is the concentration of iodine, I₂, molecules in a solution containing 2.54 g of iodine in 250 cm³ of solution?


A 0.01 mol/dm³ B 0.02 mol/dm³ C 0.04 mol/dm³ D 0.08 mol/dm³

- 18 A solution of hydrochloric acid has a concentration of 2 mol/dm^3 .

Different volumes of the acid are added to different volumes of aqueous sodium hydroxide.




The maximum temperature of each mixture is measured. The graph shows the results.



What is the concentration of the aqueous sodium hydroxide?

- A 0.67 mol/dm^3
- B 1.3 mol/dm^3
- C 1.5 mol/dm^3
- D 3.0 mol/dm^3

- 19 The diagram shows apparatus for measuring the volume of hydrogen given off when an excess of dilute hydrochloric acid is added to powdered metal. The volume of gas is measured at room temperature and pressure.



The experiment is carried out three times, using the same mass of powder each time but with different powders:

- pure magnesium
- pure zinc
- a mixture of magnesium and zinc

Which powder gives the greatest volume of hydrogen and which the least volume?

	greatest volume of H_2	least volume of H_2
A	magnesium	zinc
B	magnesium	the mixture
C	zinc	magnesium
D	zinc	the mixture

- 20 The following statements about dilute sulfuric acid are all correct.

- 1 Addition of Universal Indicator shows that the solution has a pH value of less than 7.0.
- 2 A white precipitate is formed when aqueous barium chloride is added.
- 3 The solution reacts with copper(II) oxide, forming a blue solution.
- 4 When electrolysed, hydrogen and oxygen gases are produced.

Which two statements confirm the acidic nature of the solution?

- A** 1 and 2 **B** 1 and 3 **C** 2 and 4 **D** 3 and 4

21 A solution of W has the following properties.

- When added in excess to solid ammonium chloride, a gas is given off that turns damp red litmus paper blue.
- When added in excess to a solution of pH 3, the resulting solution has a pH of 13.

What is W?


- A a strong acid
- B a strong base
- C a weak acid
- D a weak base

22 Which equation describes the most suitable reaction for making lead(II) sulfate?

- A $\text{Pb} + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + \text{H}_2$
- B $\text{PbCO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + \text{CO}_2 + \text{H}_2\text{O}$
- C $\text{Pb}(\text{NO}_3)_2 + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + 2\text{HNO}_3$
- D $\text{Pb}(\text{OH})_2 + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + 2\text{H}_2\text{O}$

23 Excess marble chips were reacted with dilute hydrochloric acid in an experiment. The volume of gas produced was measured at regular time intervals.

The results obtained are plotted in the graph as follows.



The speed of reaction decreases throughout the reaction until it comes to a stop.

Which of the following explains why the speed of reaction decreases?

- A Marble chips were completely used up.
- B Hydrochloric acid was completely used up.
- C The mass of marble chips decreased.
- D The concentration of hydrochloric acid decreased.

24 The following statements describe how the speed of reactions can be increased.

- 1 increase the amount of kinetic energy reactant particles can possess
- 2 increase the frequency of effective collisions
- 3 lower the activation energy of the reaction


Which statements describe the effect when temperature is increased?

- A 1 and 2 B 1 and 3 C 2 and 3 D 1, 2 and 3

25 In experiment 1, an excess of finely powdered marble is added to 20 cm³ of dilute hydrochloric acid.

In experiment 2, carried out under the same conditions of temperature and pressure, an excess of marble chips is added to 20 cm³ of dilute hydrochloric acid of the same concentration.


The total volumes of carbon dioxide given off are determined at intervals and plotted against time.



Which pair of curves is obtained in the two experiments?

	experiment 1	experiment 2
A	X	Z
B	X	Y
C	Y	Z
D	Y	X

- 26 The diagram shows apparatus used to investigate the conductivity of different solutions.




Which substance, in aqueous solution of concentration 1 mol/dm^3 , would cause the lamp to give the brightest light?

- A ammonia
 - B ethanoic acid
 - C dilute sodium chloride solution
 - D sulfuric acid
- 27 The heat-reflecting shields of some space rockets are gold-plated, using electrolysis.

Which electrodes and electrolyte would be used to gold-plate the heat shield?

	negative electrode	positive electrode	electrolyte
A	carbon	heat shield	gold compound
B	gold	heat shield	copper compound
C	heat shield	carbon	copper compound
D	heat shield	gold	gold compound


- 28 The following set-up shows how a block of impure copper can be purified.



The loss in mass of the anode is 50 g and the gain in mass of the cathode is 45 g.

What is the percentage purity of this sample of copper?


- A 10.0 % B 11.1 % C 90.0 % D 95.0 %
- 29 Two cells were set up as shown in the diagram. The arrow shows the direction of electron flow in the external circuit.



Which set of metals would give the electron flows in the direction shown?


	metal X	metal Y	metal Z
A	Ag	Cu	Zn
B	Ag	Zn	Cu
C	Cu	Zn	Ag
D	Zn	Cu	Ag

- 30 The energy diagram for the reaction between sodium hydroxide and hydrochloric acid is shown.



Which quantity of heat is liberated when 100 cm³ of 1 mol/dm³ hydrochloric acid reacts with 100 cm³ of 1 mol/dm³ sodium hydroxide?

- A 0.54 kJ B 2.70 kJ C 5.40 kJ D 10.8 kJ
- 31 The diagram shows the energy profile for a reaction.



Which statements about this reaction are correct?

- 1 More energy is absorbed to break the bonds than is released when new bonds are formed.
- 2 Route 1 and route 2 give the same overall equation for the reaction.
- 3 Route 2 involves the use of a catalyst.
- 4 The reaction is exothermic.

- A 1, 2 and 3 B 1 and 2 only C 2, 3 and 4 D 3 and 4 only

- 32 The equation shows a reversible reaction.



The forward reaction is endothermic.


Which of these changes will increase the yield of NO_2 ?

	pressure	temperature
A	decreased	decreased
B	decreased	increased
C	increased	decreased
D	increased	increased

- 33 Which of the following is not an example of oxidation?

- A converting iron(III) salts to iron(II) salts
- B converting magnesium atoms into magnesium ions
- C dissolving a copper anode during electrolysis
- D liberating chlorine from a chloride solution

- 34 Sulfur dioxide reacts with aqueous bromine according to the following equation.



Which element has been oxidised?


- A bromine
- B hydrogen
- C oxygen
- D sulfur

- 35 Some metals and the compounds in their ores are shown.

metal	Al	Ca	Pb	Na	Fe	Mg
compound in their ore	Al_2O_3	CaCO_3	PbS	NaCl	Fe_2O_3	MgCO_3

Which type of reaction occurs in the extraction of all of these metals from their respective ores?

- A decomposition by heat
 B electrolysis
 C precipitation
 D reduction
- 36 The diagram shows the apparatus in an experiment to reduce substance Q with the gas generated in the flask.



What are substances P and Q?

	P	Q
A	copper	copper(II) oxide
B	lead	lead(II) oxide
C	magnesium	zinc oxide
D	zinc	copper(II) oxide

- 37 Brass is an alloy.

Which statement about brass is correct?

- A It contains a sea of electrons.
- B It contains positive and negative ions which are free to move.
- C It is a compound of a metal and a non-metal.
- D It is a compound of two or more metals.

- 38 Iron is extracted from its ore haematite, Fe_2O_3 , by a reduction process in the blast furnace.

Which equation for reactions in the blast furnace shows the formation of the reducing agent?



- A $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$
- B $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$
- C $\text{CO}_2 + \text{C} \rightarrow 2\text{CO}$
- D $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$

- 39 Cars have catalytic converters fitted to reduce problems caused by some of the exhaust gases. However, cars fitted with catalytic converters still give out environmentally harmful gases.

Which of the following correctly states the harmful gas and the problem the gas causes?

	harmful gas	problem
A	nitrogen dioxide	dissolves in rain to corrode marble buildings
B	nitrogen dioxide	causes breathing problems when inhaled.
C	carbon dioxide	binds with haemoglobin in blood causing respiratory problems
D	carbon dioxide	causes the greenhouse effect leading to global warming

- 40 Nitrogen monoxide, NO, damages the ozone layer by reacting with ozone in a two-step reaction.



One nitrogen monoxide molecule can destroy thousands of ozone molecules.

Which statement correctly explains why?

- A Nitrogen monoxide in step 1 is easily generated through thunderstorms.
- B Nitrogen monoxide, while is used up in step 1, is regenerated in step 2.
- C Nitrogen dioxide produced can dissolve in rain to react with thousands of ozone molecules.
- D Nitrogen monoxide can react continuously with ozone since the ozone layer consists of thousands of ozone molecules.

End of Paper

The Periodic Table of Element

Group																				
I	II	Key										III	IV	V	VI	VII	0			
		proton (atomic) number atomic symbol name relative atomic mass																		
3 Li lithium 7	4 Be beryllium 9											1 H hydrogen 1								2 He helium 4
11 Na sodium 23	12 Mg magnesium 24																		9 F fluorine 19	
												5 B boron 11	6 C carbon 12	7 N nitrogen 14	8 O oxygen 16	17 Cl chlorine 35.5	18 Ar argon 40			
												13 Al aluminium 27	14 Si silicon 28	15 P phosphorus 31	16 S sulfur 32	19 K potassium 39				
												31 Ga gallium 70	32 Ge germanium 73	33 As arsenic 75	34 Se selenium 79	35 Br bromine 80	36 Kr krypton 84			
												49 In indium 115	50 Sn tin 119	51 Sb antimony 122	52 Te tellurium 128	53 I iodine 127	54 Xe xenon 131			
												81 Tl thallium 204	82 Pb lead 207	83 Bi bismuth 209	84 Po polonium —	85 At astatine —	86 Rn radon —			
																			116 Lv livermorium —	
												112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —			
												118 Og oganeson —	119 Uue unbinilium —	120 Uuh untrium —	121 Uut ununilium —	122 Uuq ununium —	123 Uub unbium —			
												124 Uut ununium —	125 Uuq ununium —	126 Uub unbium —	127 Uut ununium —	128 Uuq ununium —	129 Uub unbium —			
												130 Uut ununium —	131 Uuq ununium —	132 Uub unbium —	133 Uut ununium —	134 Uuq ununium —	135 Uub unbium —			
												136 Uut ununium —	137 Uuq ununium —	138 Uub unbium —	139 Uut ununium —	140 Uuq ununium —	141 Uub unbium —			
												142 Uut ununium —	143 Uuq ununium —	144 Uub unbium —	145 Uut ununium —	146 Uuq ununium —	147 Uub unbium —			
												148 Uut ununium —	149 Uuq ununium —	150 Uub unbium —	151 Uut ununium —	152 Uuq ununium —	153 Uub unbium —			
												154 Uut ununium —	155 Uuq ununium —	156 Uub unbium —	157 Uut ununium —	158 Uuq ununium —	159 Uub unbium —			
												160 Uut ununium —	161 Uuq ununium —	162 Uub unbium —	163 Uut ununium —	164 Uuq ununium —	165 Uub unbium —			
												166 Uut ununium —	167 Uuq ununium —	168 Uub unbium —	169 Uut ununium —	170 Uuq ununium —	171 Uub unbium —			
												172 Uut ununium —	173 Uuq ununium —	174 Uub unbium —	175 Uut ununium —	176 Uuq ununium —	177 Uub unbium —			
												178 Uut ununium —	179 Uuq ununium —	180 Uub unbium —	181 Uut ununium —	182 Uuq ununium —	183 Uub unbium —			
												184 Uut ununium —	185 Uuq ununium —	186 Uub unbium —	187 Uut ununium —	188 Uuq ununium —	189 Uub unbium —			
												190 Uut ununium —	191 Uuq ununium —	192 Uub unbium —	193 Uut ununium —	194 Uuq ununium —	195 Uub unbium —			
												196 Uut ununium —	197 Uuq ununium —	198 Uub unbium —	199 Uut ununium —	200 Uuq ununium —	201 Uub unbium —			
												202 Uut ununium —	203 Uuq ununium —	204 Uub unbium —	205 Uut ununium —	206 Uuq ununium —	207 Uub unbium —			
												208 Uut ununium —	209 Uuq ununium —	210 Uub unbium —	211 Uut ununium —	212 Uuq ununium —	213 Uub unbium —			
												214 Uut ununium —	215 Uuq ununium —	216 Uub unbium —	217 Uut ununium —	218 Uuq ununium —	219 Uub unbium —			
												220 Uut ununium —	221 Uuq ununium —	222 Uub unbium —	223 Uut ununium —	224 Uuq ununium —	225 Uub unbium —			
												226 Uut ununium —	227 Uuq ununium —	228 Uub unbium —	229 Uut ununium —	230 Uuq ununium —	231 Uub unbium —			
												232 Uut ununium —	233 Uuq ununium —	234 Uub unbium —	235 Uut ununium —	236 Uuq ununium —	237 Uub unbium —			
												238 Uut ununium —	239 Uuq ununium —	240 Uub unbium —	241 Uut ununium —	242 Uuq ununium —	243 Uub unbium —			
												244 Uut ununium —	245 Uuq ununium —	246 Uub unbium —	247 Uut ununium —	248 Uuq ununium —	249 Uub unbium —			
												250 Uut ununium —	251 Uuq ununium —	252 Uub unbium —	253 Uut ununium —	254 Uuq ununium —	255 Uub unbium —			
												256 Uut ununium —	257 Uuq ununium —	258 Uub unbium —	259 Uut ununium —	260 Uuq ununium —	261 Uub unbium —			
												262 Uut ununium —	263 Uuq ununium —	264 Uub unbium —	265 Uut ununium —	266 Uuq ununium —	267 Uub unbium —			
												268 Uut ununium —	269 Uuq ununium —	270 Uub unbium —	271 Uut ununium —	272 Uuq ununium —	273 Uub unbium —			
												274 Uut ununium —	275 Uuq ununium —	276 Uub unbium —	277 Uut ununium —	278 Uuq ununium —	279 Uub unbium —			
												280 Uut ununium —	281 Uuq ununium —	282 Uub unbium —	283 Uut ununium —	284 Uuq ununium —	285 Uub unbium —			
												286 Uut ununium —	287 Uuq ununium —	288 Uub unbium —	289 Uut ununium —	290 Uuq ununium —	291 Uub unbium —			
												292 Uut ununium —	293 Uuq ununium —	294 Uub unbium —	295 Uut ununium —	296 Uuq ununium —	297 Uub unbium —			
												298 Uut ununium —	299 Uuq ununium —	300 Uub unbium —	301 Uut ununium —	302 Uuq ununium —	303 Uub unbium —			
												304 Uut ununium —	305 Uuq ununium —	306 Uub unbium —	307 Uut ununium —	308 Uuq ununium —	309 Uub unbium —			
												310 Uut ununium —	311 Uuq ununium —	312 Uub unbium —	313 Uut ununium —	314 Uuq ununium —	315 Uub unbium —			
												316 Uut ununium —	317 Uuq ununium —	318 Uub unbium —	319 Uut ununium —	320 Uuq ununium —	321 Uub unbium —			
												322 Uut ununium —	323 Uuq ununium —	324 Uub unbium —	325 Uut ununium —	326 Uuq ununium —	327 Uub unbium —			
												328 Uut ununium —	329 Uuq ununium —	330 Uub unbium —	331 Uut ununium —	332 Uuq ununium —	333 Uub unbium —			
												334 Uut ununium —	335 Uuq ununium —	336 Uub unbium —	337 Uut ununium —	338 Uuq ununium —	339 Uub unbium —			
												340 Uut ununium —	341 Uuq ununium —	342 Uub unbium —	343 Uut ununium —	344 Uuq ununium —	345 Uub unbium —			
												346 Uut ununium —	347 Uuq ununium —	348 Uub unbium —	349 Uut ununium —	350 Uuq ununium —	351 Uub unbium —			
												352 Uut ununium —	353 Uuq ununium —	354 Uub unbium —	355 Uut ununium —	356 Uuq ununium —	357 Uub unbium —			
												358 Uut ununium —	359 Uuq ununium —	360 Uub unbium —	361 Uut ununium —	362 Uuq ununium —	363 Uub unbium —			
												364 Uut ununium —	365 Uuq ununium —	366 Uub unbium —	367 Uut ununium —	368 Uuq ununium —	369 Uub unbium —			
												370 Uut ununium —	371 Uuq ununium —	372 Uub unbium —	373 Uut ununium —	374 Uuq ununium —	375 Uub unbium —			
												376 Uut ununium —	377 Uuq ununium —	378 Uub unbium —	379 Uut ununium —	380 Uuq ununium —	381 Uub unbium —			
												382 Uut ununium —	383 Uuq ununium —	384 Uub unbium —	385 Uut ununium —	386 Uuq ununium —	387 Uub unbium —			
												388 Uut ununium —	389 Uuq ununium —	390 Uub unbium —	391 Uut ununium —	392 Uuq ununium —	393 Uub unbium —			
												394 Uut ununium —	395 Uuq ununium —	396 Uub unbium —	397 Uut ununium —	398 Uuq ununium —	399 Uub unbium —			
												400 Uut ununium —	401 Uuq ununium —	402 Uub unbium —	403 Uut ununium —	404 Uuq ununium —	405 Uub unbium —			
												406 Uut ununium —	407 Uuq ununium —	408 Uub unbium —	409 Uut ununium —	410 Uuq ununium —	411 Uub unbium —			
												412 Uut ununium —	413 Uuq ununium —	414 Uub unbium —	415 Uut ununium —	416 Uuq ununium —	417 Uub unbium —			
												418 Uut ununium —	419 Uuq ununium —	420 Uub unbium —	421 Uut ununium —	422 Uuq ununium —	423 Uub unbium —			
												424 Uut ununium —	425 Uuq ununium —	426 Uub unbium —	427 Uut ununium —	428 Uuq ununium —	429 Uub unbium —			
												430 Uut ununium —	431 Uuq ununium —	432 Uub unbium —	433 Uut ununium —	434 Uuq ununium —	435 Uub unbium —			
												436 Uut ununium —	437 Uuq ununium —	438 Uub unbium —	439 Uut ununium —	440 Uuq ununium —	441 Uub unbium —			
												442 Uut ununium —	443 Uuq ununium —	444 Uub unbium —	445 Uut ununium —	446 Uuq ununium —	447 Uub unbium —			
												448 Uut ununium —	449 Uuq ununium —	450 Uub unbium —	451 Uut ununium —	452 Uuq ununium —	453 Uub unbium —			
												454 Uut ununium —	455 Uuq ununium —	456 Uub unbium —	457 Uut ununium —	458 Uuq ununium —	459 Uub unbium —			
												460 Uut ununium —	461 Uuq ununium —	462 Uub unbium —	463 Uut ununium —	464 Uuq ununium —	465 Uub unbium —			
												466 Uut ununium —	467 Uuq ununium —	468 Uub unbium —	469 Uut ununium —	470 Uuq ununium —	471 Uub unbium —			
												472 Uut ununium —	473 Uuq ununium —	474 Uub unbium —	475 Uut ununium —	476 Uuq ununium —	477 Uub unbium —			
												478 Uut ununium —	479 Uuq ununium —	480 Uub unbium —	481 Uut ununium —	482 Uuq ununium —	483 Uub unbium —			
												484 Uut ununium —	485 Uuq ununium —	486 Uub unbium —	487 Uut ununium —	488 Uuq ununium —	489 Uub unbium —			
												490 Uut ununium —	491 Uuq ununium —	492 Uub unbium —	493 Uut ununium —	494 Uuq ununium —	495 Uub unbium —			
												496 Uut ununium —	497 Uuq ununium —	498 Uub unbium —	499 Uut ununium —	500 Uuq ununium —	501 Uub unbium —			
												502 Uut ununium —	503 Uuq ununium —	504 Uub unbium —	505 Uut ununium —	506 Uuq ununium —	507 Uub unbium —			
												508 Uut ununium —	509 Uuq ununium —	510 Uub unbium —	511 Uut ununium —	512 Uuq ununium —	513 Uub unbium —			
												514 Uut ununium —	515 Uuq ununium —	516 Uub unbium —	517 Uut ununium —	518 Uuq ununium —	519 Uub unbium —			
												520 Uut ununium —	521 Uuq ununium —	522 Uub unbium —	523 Uut ununium —	524 Uuq ununium —	525 Uub unbium —			
												526 Uut ununium —	527 Uuq ununium —	528 Uub unbium —	529 Uut ununium —	530 Uuq ununium —	531 Uub unbium —			
												532 Uut ununium —	533 Uuq ununium —	534 Uub unbium —	535 Uut ununium —	536 Uuq ununium —	537 Uub unbium —			
												538 Uut ununium —	539 Uuq ununium —	540 Uub unbium —	541 Uut ununium —	542 Uuq ununium —	543 Uub unbium —			
												544 Uut ununium —	545 Uuq ununium —	546 Uub unbium —	547 Uut ununium —	548 Uuq ununium —	549 Uub unbium —			
												550 Uut ununium —	551 Uuq ununium —	552 Uub unbium —	553 Uut ununium —	554 Uuq ununium —	555 Uub unbium —			
												556 Uut ununium —	557 Uuq ununium —	558 Uub unbium —	559 Uut ununium —	560 Uuq ununium —	561 Uub unbium —			
												562 Uut ununium —	563 Uuq ununium —	564 Uub unbium —	565 Uut ununium —	566 Uuq ununium —	567 Uub unbium —			
												568 Uut ununium —	569 Uuq ununium —	570 Uub unbium —	571 Uut ununium —	572 Uuq ununium —	573 Uub unbium —			
												574 Uut ununium —	575 Uuq ununium —	576 Uub unbium —	577 Uut ununium —	578 Uuq ununium —	579 Uub unbium —			
												580 Uut ununium —	581 Uuq ununium —	582 Uub unbium —	583 Uut ununium —	584 Uuq ununium —	585 Uub unbium —			
												586 Uut ununium —	587 Uuq ununium —	588 Uub unbium —	589 Uut ununium —	590 Uuq ununium —	591 Uub unbium —			
												592 Uut ununium —	593 Uuq ununium —	594 Uub unbium —	595 Uut ununium —	596 Uuq ununium —	597 Uub unbium —			
												598 Uut ununium —	599 Uuq ununium —	600 Uub unbium —	601 Uut ununium —	602 Uuq ununium —	603 Uub unbium —			
												604 Uut ununium —	605 Uuq ununium —	606 Uub unbium —	607 Uut ununium —	608 Uuq ununium —	609 Uub unbium —			
												610 Uut ununium —	611 Uuq ununium —							

21

Sc

scandium

45

39

Y

yttrium

89

57 – 71

lanthanoids

—

89 – 103

actinoids

—

22

Ti

titanium

48

40

Zr

zirconium

91

72

Hf

hafnium

178

104

Rf

Rutherfordium

—

23

V

vanadium

51

41

Nb

niobium

93

73

Ta

tantalum

181

105

Db

dubnium

—

24

Cr

chromium

52

42

Mo

molybdenum

96

74

W

tungsten

184

106

Sg

seaborgium

—

25

Mn

manganese

55

43

Tc

technetium

—

75

Re

rhenium

186

107

Bh

bohrium

—

26

Fe

iron

56

44

Ru

ruthenium

101

76

Os

osmium

190

108

Hs

hassium

—

27

Co

cobalt

59

45

Rh

rhodium

103

77

Ir

iridium

192

109

Mt

meitnerium

—

28

Ni

nickel

59

46

Pd

palladium

106

78

Pt

platinum

195

110

Ds

darmstadtium

—

29

Cu

copper

64

47

Ag

silver

108

79

Au

gold

197

111

Rg

roentgenium

—

30

Zn

zinc

65

48

Cd

cadmium

112

80

Hg

mercury

201

112

Cn

copernicium

—

31

Ga

gallium

70

49

In

indium

115

81

Tl

thallium

204

114

Fl

flerovium

—

32

Ge

germanium

73

50

Sn

tin

119

82

Pb

lead

207

116

Lv

livermorium

—

33

As

arsenic

75

51

Sb

antimony

122

83

Bi

bismuth

209

116

Lv

livermorium

—

34

Se

selenium

79

52

Te

tellurium

128

84

Po

polonium

—

116

Lv

livermorium

—

35

Br

bromine

80

53

I

iodine

127

85

At

astatine

—

116

Lv

livermorium

—

36

Kr

krypton

84

54

Xe

xenon

131

86

Rn

radon

—

116

Lv

livermorium

—

37

Br

bromine

80

53

I

iodine

127

85

At

astatine

—

116

Lv

livermorium

—

38

Sr

strontium

88

54

Xe

xenon

131

86

Rn

radon

—

116

Lv

livermorium

—

39

Y

yttrium

89

55

Cs

caesium

133

87

Fr

francium

—

116

Lv

livermorium

—

40

Ca

calcium

40

56

Ba

barium

137

88

Ra

radium

—

116

Lv

livermorium

—

41

Nb

niobium

93

57

Cs

caesium

133

87

Fr

francium

—

116

Lv

livermorium

—

42

Mo

molybdenum

96

58

Ba

barium

137

88

Ra

radium

—

116

Lv

livermorium

—

43

Tc

technetium

—

59

Cs

caesium

133

89

Fr

francium

—

116

Lv

livermorium

—

44

Ru

ruthenium

101

60

Ba

barium

137

90

Th

thorium

232

116

Lv

livermorium

—

45

Rh

rhodium

103

61

Pm

promethium

—

91

Pa

protactinium

231

116

Lv

livermorium

—

46

Pd

palladium

106

62

Sm

samarium

150

92

U

uranium

238

116

Lv

livermorium

—

47

Ag

silver

108

63

Eu

europlum

152

93

Np

neptunium

—

116

Lv

livermorium

—

48

Cd

cadmium

112

64

Gd

gadolinium

157

94

Pu

plutonium

—

116

Lv

livermorium

—

49

In

indium

115

65

Tb

terbium

159

95

Am

americium

—

116

Lv

livermorium

—

50

Sn

tin

119

66

Dy

dysprosium

163

96

Cm

curium

—

116

Lv

livermorium

—

51

Sb

antimony

122

67

Ho

holmium

165

97

Bk

berkelium

—

116

Lv

livermorium

—

52

Te

tellurium

128

68

Er

erbium

167

98

Cf

californium

—

116

Lv

livermorium

—

53

I

iodine

127

69

Tm

thulium

169

99

Es

einsteinium

—

116

Lv

livermorium

—

54

Xe

xenon

131

70

Yb

ytterbium

173

100

Fm

fermium

—

116

Lv

livermorium

—

55

Cs

caesium

133

71

Lu

lutetium

175

101

Md

mandeleevium

—

116

Lv

livermorium

—

56

Ba

barium

137

72

Hf

hafnium

178

102

No

nobelium

—

116

Lv

livermorium

—

57

La

lanthanum

139

73

Ta

tantalum

181

103

Lr

lawrencium

—

116

Lv

livermorium

—

58

Ce

cerium

140

74

W

tungsten

184

104

Rf

Rutherfordium

—

116

Lv

livermorium

—

59

Pr

praseodymium

141

75

Re

rhenium

186

105

Db

dubnium

—

116

Lv

livermorium

—

60

Nd

neodymium

144

76

Os

osmium

190

106

Sg

seaborgium

—

116

Lv

livermorium

—

61

Pm

promethium

—

77

Ir

iridium

192

107

Bh

bohrium

—

116

Lv

livermorium

—

62

Sm

samarium

150

78

Pt

platinum

195

108

Hs

hassium

—

116

Lv

livermorium

—

63

Eu

europlum

152

79

Au

gold

197

109

Mt

meitnerium

—

116

Lv

livermorium

—

64

Gd

gadolinium

157

80

Hg

mercury

201

110

Ds

darmstadtium

—

116

Lv

livermorium

—

65

Tb

terbium

159

81

Tl

thallium

204

111

Rg

roentgenium

—

116

Lv

livermorium

—

66

Dy

dysprosium

163

82

Pb

lead

207

112

Cn

copernicium

—

116

Lv

livermorium

—

67

Ho

holmium

165

83

Bi

bismuth

209

113

Nh

nihonium

—

116

Lv

livermorium

—

68

Er

erbium

167

84

Cd

cadmium

112

114

Fl

flerovium

—

116

Lv

livermorium

—

69

Tm

thulium

169

85

Ag

silver

108

115

Mc

moscovium

—

116

Lv

livermorium

—

70

Yb

ytterbium

173

86

Rn

radon

—

116

Lv

livermorium

—

116

Lv

livermorium

—

71

Lu

lutetium

175

87

Fr

francium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

72

Hf

hafnium

178

88

Ra

radium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

73

Ta

tantalum

181

89

Ac

actinium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

74

W

tungsten

184

90

Th

thorium

232

116

Lv

livermorium

—

116

Lv

livermorium

—

75

Re

rhenium

186

91

Pa

protactinium

231

116

Lv

livermorium

—

116

Lv

livermorium

—

76

Os

osmium

190

92

U

uranium

238

116

Lv

livermorium

—

116

Lv

livermorium

—

77

Ir

iridium

192

93

Np

neptunium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

78

Pt

platinum

195

94

Pu

plutonium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

79

Au

gold

197

95

Am

americium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

80

Hg

mercury

201

96

Cm

curium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

81

Tl

thallium

204

97

Bk

berkelium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

82

Pb

lead

207

98

Cf

californium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

83

Bi

bismuth

209

99

Es

einsteinium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

84

Po

polonium

—

100

Fm

fermium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

85

At

astatine

—

101

Md

mandeleevium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

86

Rn

radon

—

102

No

nobelium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

87

Fr

francium

—

103

Lr

lawrencium

—

116

Lv

livermorium

—

116

Lv

livermorium

—

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).



**JUNYUAN SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2020
SECONDARY FOUR EXPRESS**

CANDIDATE NAME

--

CLASS

4	E	
---	---	--

INDEX NUMBER

--	--

CHEMISTRY

6092 / 02

Paper 2

24 August 2020

1 hour 45 minutes

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in.
Write in dark blue or black pen on both sides of the paper.
You may use a soft pencil for any diagrams, graphs or rough workings.
Do not use paper clips, highlighters, glue or correction fluid.

The use of an approved scientific calculator is expected, where appropriate.

Section A

Answer all the questions in the spaces provided.

Section B

Answer all **three** questions, the last question is in the form either/or.
Write your answers in the spaces provided.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [] at the end of each question or part question.
A copy of the Periodic Table is printed on page 23.

For Examiner's Use	
Section A	
Section B	
Total	80

This document consists of 23 printed pages

[Turn Over

Section A

Answer all questions in this section in the spaces provided. The total mark for this section is 50.

A1 Use the list of substances to answer the questions.


ammonia	iodine	copper
hydrochloric acid	calcium chloride	nitrogen
argon	sulfur dioxide	magnesium

Which substance

- (a) reacts with gaseous hydrogen chloride to give a white solid?
[1]
- (b) changes directly into a gas when gently heated?
[1]
- (c) displaces iron from its salt solution?
[1]
- (d) exists as a monatomic gas?
[1]
- (e) is a colourless solution which can be used to distinguish between aqueous silver nitrate and aqueous zinc nitrate?
[1]

[Total: 5]

- A2** A sample of black ink contains a mixture of red, blue and yellow dyes. Usually, the solvent used to separate the dyes in black ink is a mixture of ethanol and water. The coloured dyes have different R_f values in solvents with different proportions of ethanol in the mixture as shown in the graph.



- (a) Deduce the R_f value of the blue dye on the chromatogram when the solvent is a mixture of 32 cm³ of ethanol and 168 cm³ of water.

R_f value [1]

- (b) Based on the graph, justify whether a pure solvent of water is suitable for the separation of the black ink using paper chromatography.

.....

 [2]

- (c) A student carried out chromatography on the black ink using a mixture of ethanol and water as the solvent. He discovered only one spot forming on the resulting chromatogram.

Using evidence from the graph, explain why he cannot conclude that the black ink is a pure substance.

.....

 [2]


- (d) Draw a labelled set up for the separation above.

[2]

[Total: 7]

- A3** Elemental phosphorus exists as two forms – white and black phosphorus. The different forms display strikingly different properties.

The table shows the structures and properties of white and black phosphorus.

	white phosphorus	black phosphorus
structure		
properties at room temperature	waxy white solid	black crystalline solid with a greasy touch
melting point / °C	44.2	610

- (a) Give one similarity and one difference in the bonding and structure between the two forms of phosphorus.

similarity:

.....

difference:

.....[2]

6092/PRELIM/4E/2020

- (b) Explain, in terms of bonding, why the melting point of black phosphorus is much higher than the melting point of white phosphorus.

.....

.....

.....

.....

.....[3]

- (c) White phosphorus reacts with chlorine to form phosphorus chloride, PCl_3 . Draw a 'dot-and-cross' diagram to show the bonding in a molecule of PCl_3 .

Show only valence electrons.

[2]

[Total: 7]

- A4** The relative positions of the elements rubidium (Rb), beryllium (Be) and bismuth (Bi) in the reactivity series are shown in the table below.

position in the reactivity series (highest to lowest)
rubidium
sodium
magnesium
beryllium
iron
hydrogen
bismuth
copper
silver

You may assume that these elements do not show variable valencies.

- (a) An unknown photograph showing specks of silvery deposits with the caption - "Pure rubidium found on a tiny island in the Pacific Ocean" was posted in the early morning of August 16, 2016, on social media. The post has since gone viral.

Using the information above and your knowledge of metals in the reactivity series, discuss the validity of this post.

.....

[2]

- (b) Predict, with reasons, the reactions of beryllium with cold water and steam.

.....

[4]


- (c) Suggest a suitable method to extract bismuth from its ore.

.....[1]


[Total: 7]

6092/PRELIM/4E/2020

- A5** Barium hydroxide reacts with sulfuric acid to form barium sulfate precipitate. The equation for this reaction is as shown.



In an experiment, 0.500 mol/dm^3 sulfuric acid was gradually added to 25.0 cm^3 of barium hydroxide solution in a conical flask. The mixture was continuously stirred with an iron stirrer covered in plastic. The mixture was connected to an ammeter. The reading was taken after the addition of every 1.00 cm^3 of sulfuric acid. The graph below shows the results obtained from the experiment.



- (a) Ignoring the ions contributed by the ionisation of water, state the formulae of the ions present in the conical flask when the following volumes of sulfuric acid were added to the barium hydroxide solution:

(i) 0.00 cm^3

(ii) 8.00 cm^3 [2]

- (b) Explain the change in ammeter reading from the start of the experiment to the point when 5.00 cm^3 of acid was added.

.....

 [3]

(c) Calculate the concentration, in mol/dm^3 , of barium hydroxide solution used.

[2]

(d) Suggest why the iron stirrer was covered in plastic.


.....

.....[1]

[Total: 8]

A6 Solid P is a metal carbonate.

The graph shows how the mass of the solid in a boiling tube changes as the experiment progresses.



There are four stages for the experiment.

stages	description
I	solid P is heated strongly with a non-luminous flame to form solid Q.
II	excess dilute hydrochloric acid is added to solid Q.
III	aqueous ammonia is added to the reaction mixture from stage II.
IV	aqueous silver nitrate is added to the reaction mixture from stage III.

(a) Suggest a possible identity of the cation present in solid P.

Explain your answer.

.....

[3]

- (b) Do you expect any carbon dioxide to be evolved in stage II?

Explain your answer.

.....
[1]


- (c) Write the ionic equation for the formation of the solid in stage IV.

.....[2]


[Total: 6]

- A7** An experiment was set up to determine the percentage of oxygen in air.

Two gas syringes were connected to a small tube containing excess iron, as shown in the diagram below.



At the start of the experiment, the apparatus contained a total of 200 cm³ of air. During heating, the iron reacted with oxygen in the air to form black iron(II) oxide.



The iron was heated until the volume of gas, measured at room temperature and pressure, remained constant.

- (a) Explain why it was important to continue heating until the volume of gas remained constant.

.....[1]


- (b) The volume of gas left after the reaction was complete was 160 cm³.

Calculate the percentage of oxygen in the air.

[1]

6092/PRELIM/4E/2020

- (c) Iron(II) oxide can be obtained by the thermal decomposition of iron(II) oxalate.



- (i) Determine the oxidation states of carbon in the following substances.

	oxidation state of carbon
FeC_2O_4	
CO_2	
CO	

[2]

- (ii) Explain, in terms of oxidation states, why the above reaction is a redox reaction.


.....

.....


.....[2]

[Total: 6]

- A8** Ammonia is prepared industrially from hydrogen and nitrogen in the presence of a suitable catalyst according to the equation below.



The graph below shows the variation of the equilibrium yield of ammonia with pressure at different temperatures.



- (a) A particular industrial plant uses a pressure of 400 atm and a temperature of 500 °C. From the graph, determine the percentage yield of ammonia under these conditions.

.....[1]

- (b) The gases leaving the reactor contain unreacted nitrogen and hydrogen, and about 15% ammonia by volume. Unreacted nitrogen and hydrogen are fed back into the reactor.

Give two reasons why the unreacted gases are fed back into the reactor.

.....

[2]

- (c) Temperatures less than 400 °C are not used for this industrial reaction even though such temperatures give a greater equilibrium yield of ammonia.

Suggest why this is so.

.....

.....[1]

[Total: 4]

SECTION B

Answer all three questions from this section.

The last question is in the form of either/or and only one of the alternatives should be attempted.

The total mark for this section is 30.

- B9** A fuel cell is a chemical cell in which reactants are continuously supplied to produce electricity.


Two such cells are the **Alkaline Fuel Cell (AFC)** and the **Proton Exchange Membrane Fuel Cell (PEMFC)**

Alkaline Fuel Cell

AFCs use an alkaline electrolyte such as potassium hydroxide in water and are generally fuelled with pure hydrogen. Typical operating temperatures are around 70 °C. As a result of the low operating temperature, a variety of non-precious metals can be used as catalysts to speed up the reactions occurring at the anode and cathode.

At the anode, the hydrogen gas reacts with the hydroxide ions to form water. The water then travels through the membrane to the cathode side of the cell where they then react with oxygen to form hydroxide ions. The electrons travel in an external circuit, generating the electrical output of the cell.

AFC cell type is easily poisoned by carbon dioxide (CO_2). In fact, even the small amount of CO_2 in the air can affect this cell's operation, making it necessary to purify both the hydrogen and oxygen used in the cell. This purification process is costly.




6092/PRELIM/4E/2020

Proton Exchange Membrane Fuel Cell

The **PEMFC** uses a water-based, acidic polymer membrane as its electrolyte, with platinum-based electrodes. PEMFC cells operate at relatively low temperatures (below 100 °C). Due to the use of precious metal-based electrodes, these cells must operate on pure hydrogen.

Hydrogen gas is processed at the anode where electrons are separated to form hydrogen ions on the surface of a platinum-based catalyst. The hydrogen ions pass through the membrane to the cathode side of the cell where they then react with oxygen to form water. The electrons travel in an external circuit, generating the electrical output of the cell



- (a) Compare the reactions at the electrodes for **AFC** and **PEMFC**. Complete the table below with the relevant half-equations.

	cathode	anode
AFC		
PEMFC		

[4]

- (b) Write the overall equation, with state symbols, for both cells.

.....[2]

- (c) Explain, with reference to the nature of CO_2 , why it poisons **AFC**. Write an equation to support your answer.

.....
.....
.....[2]

- (d) Suggest one reason why the operation of **AFC** is more economical than that of **PEMFC**.

.....
.....[1]

- (e) Most of the hydrogen produced today is made via steam-methane reforming. In this process, high temperature steam reacts with methane (CH_4), in the presence of a catalyst to produce hydrogen and carbon monoxide.

- (i) Hydrogen fuel cells are environmentally-friendly.

Explain why.

.....[1]


- (ii) Suggest why some environmentalists argue against the use of hydrogen fuel cells.

.....
.....[1]

[Total: 11]

- B10** Rocket belts, also known as jetpacks, can be used to lift people into the air and transport them over short distances.

The picture below shows a rocket belt, usually strapped onto a person's back.



Rocket belts contain concentrated hydrogen peroxide solution as a fuel. An exothermic change occurs when the hydrogen peroxide decomposes rapidly to form oxygen and water.

This rapid release of oxygen from the rocket belt lifts the person off the ground.

- (a) Write a balanced chemical equation for the decomposition of hydrogen peroxide in the rocket belt.

.....[1]

- (b) Explain why the decomposition of hydrogen peroxide is exothermic, in terms of the energy changes that take place during bond breaking and bond forming.

.....
.....
.....[2]


- (c) The decomposition of hydrogen peroxide is very slow at room temperature.

In a rocket belt, silver powder is present to speed up this decomposition reaction.

The silver powder remains unchanged at the end of the reaction.

- (i) Draw an energy profile diagram below to show how the silver powder speeds up the decomposition of hydrogen peroxide.

energy (kJ)



- (ii) Explain the shape of the energy profile diagram.

[3]

.....
.....
.....
.....[2]

- (iii) Suggest another way to increase the rate of decomposition of hydrogen peroxide in the rocket belt.

.....
.....[1]

[Total: 9]

EITHER

B11 Nickel is a transition element. It is manufactured in a four-stage process from nickel(II) sulfide, NiS.

Stage 1 - nickel(II) sulfide is heated in air to form nickel(II) oxide and sulfur dioxide

Stage 2 - nickel(II) oxide is heated with carbon to give impure nickel

Stage 3 - impure nickel is reacted with carbon monoxide to make nickel tetracarbonyl, Ni(CO)_4

Stage 4 - nickel tetracarbonyl is decomposed to give pure nickel

(a) (i) Construct the balanced equation for the reaction in stage 1.

.....[1]

(ii) Calculate the volume of sulfur dioxide formed if 182 kg of nickel(II) sulfide is used.

volume of sulfur dioxide = ... dm^3 [2]

(b) In terms of structure and bonding, explain why pure nickel is a good electrical conductor.

.....

[2]

- (c) In an experiment, small amounts of three metals were added to three aqueous metal nitrate solutions. The observations are shown in the table.

	aqueous zinc nitrate	aqueous nickel(II) nitrate	aqueous copper(II) nitrate
zinc	no visible reaction	green solution turns colourless and zinc gets coated with a grey solid	blue solution turns colourless and zinc gets coated with a pink solid
nickel	?	no visible reaction	?
copper	no visible reaction	no visible reaction	no visible reaction

Predict the observations when nickel is added to separate solutions of zinc nitrate and copper(II) nitrate.

with zinc nitrate

.....

with copper(II) nitrate

.....
[3]

- (d) Explain why this four-stage process cannot be used to manufacture magnesium.

.....

[2]

[Total: 10]

OR

B11

Cleaning solutions usually contain the acid salt, sodium dihydrogen phosphate. It is known as an 'acid salt' as it can behave as both an acid and a salt. This salt can be made by reacting sodium hydroxide with phosphoric acid, H_3PO_4 .

Sodium dihydrogen phosphate contains the anion, H_2PO_4^- .

- (a) Write an equation, with state symbols, for the formation of sodium dihydrogen phosphate.

.....[2]

- (b) The table shows information about other acidic compounds.

name	pH of a 1.0 mol/dm^3 solution
phosphoric acid	4.7
sodium dihydrogen phosphate	4.5
ethanoic acid	3.8
sulfuric acid	1.0

- (i) Identify a strong acid and a weak acid.

Explain your reasoning.

.....

.....

.....

.....

.....

.....[3]

- (ii) Describe an experiment, other than measuring pH, that you could carry out to show that the named acids in (b)(i) is a strong acid and weak acid respectively.

State what measurements you would make and what results you would expect. Include in your answers two variables to be kept constant.

You may include diagrams in your answer.

.....

.....

.....

.....

.....

.....

.....

.....[5]

[Total: 10]

End of paper

6092/PRELIM/4E/2020

The Periodic Table of Elements

Group																	
I	II	Key										III	IV	V	VI	VII	0
		proton (atomic) number atomic symbol name relative atomic mass															
3 Li lithium 7	4 Be beryllium 9											5 B boron 11	6 C carbon 12	7 N nitrogen 14	8 O oxygen 16	9 F fluorine 19	10 Ne neon 20
11 Na sodium 23	12 Mg magnesium 24											13 Al aluminium 27	14 Si silicon 28	15 P phosphorus 31	16 S sulfur 32	17 Cl chlorine 35.5	18 Ar argon 40
19 K potassium 39	20 Ca calcium 40	21 Sc scandium 45	22 Ti titanium 48	23 V vanadium 51	24 Cr chromium 52	25 Mn manganese 55	26 Fe iron 56	27 Co cobalt 59	28 Ni nickel 59	29 Cu copper 64	30 Zn zinc 65	31 Ga gallium 70	32 Ge germanium 73	33 As arsenic 75	34 Se selenium 79	35 Br bromine 80	36 Kr krypton 84
37 Rb rubidium 85	38 Sr strontium 88	39 Y yttrium 89	40 Zr zirconium 91	41 Nb niobium 93	42 Mo molybdenum 96	43 Tc technetium -	44 Ru ruthenium 101	45 Rh rhodium 103	46 Pd palladium 106	47 Ag silver 108	48 Cd cadmium 112	49 In indium 115	50 Sn tin 119	51 Sb antimony 122	52 Te tellurium 128	53 I iodine 127	54 Xe xenon 131
55 Cs caesium 133	56 Ba barium 137	57-71 lanthanoids	72 Hf hafnium 178	73 Ta tantalum 181	74 W tungsten 184	75 Re rhenium 186	76 Os osmium 190	77 Ir iridium 192	78 Pt platinum 195	79 Au gold 197	80 Hg mercury 201	81 Tl thallium 204	82 Pb lead 207	83 Bi bismuth 209	84 Po polonium -	85 At astatine -	86 Rn radon -
87 Fr francium -	88 Ra radium -	89-103 actinoids	104 Rf rutherfordium -	105 Db dubnium -	106 Sg seaborgium -	107 Bh bohrium -	108 Hs hassium -	109 Mt meitnerium -	110 Ds darmstadtium -	111 Rg roentgenium -	112 Cn copernicium -	114 Fl flerovium -	116 Lv livermorium -	117 Ts tennessine -	118 Og oganesson -	119 Nh nihonium -	120 Dl dubnium -

lanthanoids														
57 La lanthanum 139	58 Ce cerium 140	59 Pr praseodymium 141	60 Nd neodymium 144	61 Pm promethium -	62 Sm samarium 150	63 Eu europium 152	64 Gd gadolinium 157	65 Tb terbium 159	66 Dy dysprosium 163	67 Ho holmium 165	68 Er erbium 167	69 Tm thulium 169	70 Yb ytterbium 173	71 Lu lutetium 175
actinoids														
89 Ac actinium -	90 Th thorium 232	91 Pa protactinium 231	92 U uranium 238	93 Np neptunium -	94 Pu plutonium -	95 Am americium -	96 Cm curium -	97 Bk berkelium -	98 Cf californium -	99 Es einsteinium -	100 Fm fermium -	101 Md mendelevium -	102 No nobelium -	103 Lr lawrencium -

The volume of one mole of any gas is 24 dm^3 at room temperature and pressure (r.t.p.).



**JUNYUAN SECONDARY SCHOOL
PRELIMINARY EXAMINATION 2020
SECONDARY FOUR EXPRESS**

CANDIDATE NAME

--

CLASS

4

E

--

INDEX NUMBER

--	--

CHEMISTRY

6092

PAPER 1

2 September 2020

1 hour

Additional Materials: Multiple Choice Answer Sheet

READ THESE INSTRUCTIONS FIRST

Write your name, class and index number on all the work you hand in.

Write in soft pencil.

Do not use paper clips, glue or correction fluid.

There are forty questions on this paper. Answer all questions. For each question there are four possible answers, A, B, C and D.

Choose the one you consider correct and record your choice in soft pencil on the separate Answer Sheet.

Read the instructions on the Answer Sheet very carefully.

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

A copy of the Periodic Table is printed on page 19.

The use of an approved scientific calculator is expected, where appropriate.

This document consists of 19 printed pages.

1 Which substance would diffuse most quickly?

- A carbon dioxide at 0 °C
- B carbon dioxide at 25 °C
- C neon at 0 °C
- D neon at 25 °C





2 A student tested a solution by adding aqueous sodium hydroxide. A precipitate was not seen because the reagent was added too quickly.

What could not have been present in the solution?

- A Al^{3+} B Ca^{2+} C NH_4^+ D Zn^{2+}


3 Four gas jars each contains one of the gases, ammonia, chlorine and oxygen. A strip of damp blue litmus paper and a strip of damp red litmus paper are placed in each jar.

In which gas jar will both the damp blue litmus paper and the damp red litmus paper change colour?



- 4 The diagram shows the chromatogram obtained by analysis of a single dye.

Three measurements are shown.



How is the R_f value of the dye calculated?

- A $\frac{x}{x+y}$ B $\frac{y}{x+y}$ C $\frac{x}{x+y+z}$ D $\frac{y}{x+y+z}$
- 5 The apparatus shown in Fig. 5.1 can be used to separate pure water from seawater.




Fig. 5.1

Two samples are taken, one at point X and another at point Y.

Which of the following statements about X and Y is incorrect?

- A X and Y can be separated into their components by physical methods.
 B When heated to dryness, X leaves a residue while Y does not.
 C X boils over a range of temperatures, while Y boils at 100 °C.
 D X is a mixture while Y is a compound.

- 6 The following statement describes substance X.

A chemical reaction takes place and heat is liberated when this white solid, X is formed.

Which of the following correctly classifies X and explains why X is classified as such?

	classification	explanation
A	element	When decomposition takes place, an element, X, is produced.
B	compound	Bond forming takes place to produce X.
C	mixture	The reactants and the product X, form a mixture.
D	either an element or a compound	A chemical reaction can produce either an element or a compound.


- 7 The atoms $^{64}_{29}\text{Cu}$ and $^{65}_{30}\text{Zn}$ have the same

- A nucleon number.
- B number of electrons.
- C number of neutrons.
- D proton number.

- 8 Which of the following correctly describes what happens when calcium atoms form calcium ions?


	calcium atoms	ionic equation for the formation
A	gain electrons	$\text{Ca} + 2\text{e}^- \rightarrow \text{Ca}^{2+}$
B	gain electrons	$\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$
C	lose electrons	$\text{Ca} + 2\text{e}^- \rightarrow \text{Ca}^{2+}$
D	lose electrons	$\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$

- 9 Element X has a lattice of positive ions and a 'sea of electrons'.




Which property will X have?

- A It conducts electricity by the movement of ions and electrons.
 B It has a high melting point.
 C It is decomposed by an electric current.
 D It is not malleable.
- 10 When a covalent substance in liquid state boils, its molecules become more widely spaced.
- Which property of the molecules has the most influence on the amount of energy required to boil a covalent substance?
- A the forces of attraction between the molecules
 B the reactivity of the molecules
 C the shape of the molecules
 D the strength of the covalent bonds in the molecules
- 11 The diagrams show the structures of two forms of carbon.



S



T

Which set of data is correct for these two structures?

	conducts electricity	very hard material	can be used as lubricant
A	T	T	S
B	S	T	S
C	S	S	T
D	T	S	T

12 Which statement about ionic compounds is correct?

- A Ionic compounds conduct electricity when solid because they contain charged particles that can move.
- B Ionic compounds consist of a lattice of positive ions and negative ions.
- C Most ionic compounds are solids at room temperature because of the strong attraction between electrons and positive ions.
- D When molten or in aqueous solution, ionic compounds conduct electricity because they contain electrons that can move.

13 Sulfur and selenium (Se) are in the same group of the Periodic Table.

From this, we would expect selenium to form compounds having the formulae

- A SeO , Na_2Se and NaSeO_4 .
- B SeO_2 , Na_2Se and NaSeO_4 .
- C SeO_2 , Na_2Se and Na_2SeO_4 .
- D SeO_3 , NaSe and NaSeO_4 .

14 Students give their own special symbols to five non-metallic elements. All five non-metals are in the same group of the Periodic Table. These non-metals exist as coloured elements.

The special symbols are shown in Fig. 14.1. The order of chemical reactivity of these non-metals is also shown.




Fig. 14.1

A solution containing Cc ions can be displaced by two of the elements in the Group.

Which of the following correctly shows the ionic equation for one such reaction?

- A $\text{Aa}(\text{aq}) + \text{Cc}^-(\text{aq}) \rightarrow \text{Aa}^-(\text{aq}) + \text{Cc}(\text{aq})$
- B $\text{Bb}_2(\text{aq}) + 2\text{Cc}^-(\text{aq}) \rightarrow 2\text{Bb}^-(\text{aq}) + \text{Cc}_2(\text{aq})$
- C $\text{Dd}(\text{aq}) + \text{Cc}^-(\text{aq}) \rightarrow \text{Dd}^-(\text{aq}) + \text{Cc}(\text{aq})$
- D $\text{Ee}_2(\text{aq}) + 2\text{Cc}^-(\text{aq}) \rightarrow 2\text{Ee}^-(\text{aq}) + \text{Cc}_2(\text{aq})$

- 15 The table shows some of the properties of four elements.

Which element is most likely to be a transition metal?

	melting point / °C	density / g/cm ³	electrical conductivity
A	3550	3.5	poor
B	1860	7.2	good
C	660	2.7	good
D	232	7.3	good

- 16 The relative formula masses of four compounds are given.

A student has a 1.0 g sample of each compound.

Which sample contains the highest number of moles of oxygen atoms?


	compound	relative formula mass
A	Al ₂ O ₃	102
B	CuO	80
C	H ₂ SO ₄	98
D	HNO ₃	63

- 17 What is the concentration of iodine, I₂, molecules in a solution containing 2.54 g of iodine in 250 cm³ of solution?


A 0.01 mol/dm³ **B** 0.02 mol/dm³ **C** 0.04 mol/dm³ **D** 0.08 mol/dm³

- 18 A solution of hydrochloric acid has a concentration of 2 mol/dm^3 .

Different volumes of the acid are added to different volumes of aqueous sodium hydroxide.




The maximum temperature of each mixture is measured. The graph shows the results.



What is the concentration of the aqueous sodium hydroxide?

- A 0.67 mol/dm^3
- B 1.3 mol/dm^3
- C 1.5 mol/dm^3
- D 3.0 mol/dm^3

- 19 The diagram shows apparatus for measuring the volume of hydrogen given off when an excess of dilute hydrochloric acid is added to powdered metal. The volume of gas is measured at room temperature and pressure.



The experiment is carried out three times, using the same mass of powder each time but with different powders:

- pure magnesium
- pure zinc
- a mixture of magnesium and zinc

Which powder gives the greatest volume of hydrogen and which the least volume?

	greatest volume of H ₂	least volume of H ₂
A	magnesium	zinc
B	magnesium	the mixture
C	zinc	magnesium
D	zinc	the mixture

- 20 The following statements about dilute sulfuric acid are all correct.

- 1 Addition of Universal Indicator shows that the solution has a pH value of less than 7.0.
- 2 A white precipitate is formed when aqueous barium chloride is added.
- 3 The solution reacts with copper(II) oxide, forming a blue solution.
- 4 When electrolysed, hydrogen and oxygen gases are produced.

Which two statements confirm the acidic nature of the solution?

- A** 1 and 2 **B** 1 and 3 **C** 2 and 4 **D** 3 and 4

21 A solution of W has the following properties.

- When added in excess to solid ammonium chloride, a gas is given off that turns damp red litmus paper blue.
- When added in excess to a solution of pH 3, the resulting solution has a pH of 13.

What is W?


- A a strong acid
- B a strong base
- C a weak acid
- D a weak base

22 Which equation describes the most suitable reaction for making lead(II) sulfate?

- A $\text{Pb} + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + \text{H}_2$
- B $\text{PbCO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + \text{CO}_2 + \text{H}_2\text{O}$
- C $\text{Pb}(\text{NO}_3)_2 + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + 2\text{HNO}_3$
- D $\text{Pb}(\text{OH})_2 + \text{H}_2\text{SO}_4 \rightarrow \text{PbSO}_4 + 2\text{H}_2\text{O}$

23 Excess marble chips were reacted with dilute hydrochloric acid in an experiment. The volume of gas produced was measured at regular time intervals.

The results obtained are plotted in the graph as follows.



The speed of reaction decreases throughout the reaction until it comes to a stop.

Which of the following explains why the speed of reaction decreases?

- A Marble chips were completely used up.
- B Hydrochloric acid was completely used up.
- C The mass of marble chips decreased.
- D The concentration of hydrochloric acid decreased.

24 The following statements describe how the speed of reactions can be increased.

- 1 increase the amount of kinetic energy reactant particles can possess
- 2 increase the frequency of effective collisions
- 3 lower the activation energy of the reaction


Which statements describe the effect when temperature is increased?

- A** 1 and 2 **B** 1 and 3 **C** 2 and 3 **D** 1, 2 and 3

25 In experiment 1, an excess of finely powdered marble is added to 20 cm³ of dilute hydrochloric acid.

In experiment 2, carried out under the same conditions of temperature and pressure, an excess of marble chips is added to 20 cm³ of dilute hydrochloric acid of the same concentration.


The total volumes of carbon dioxide given off are determined at intervals and plotted against time.



Which pair of curves is obtained in the two experiments?

	experiment 1	experiment 2
A	X	Z
B	X	Y
C	Y	Z
D	Y	X

- 26 The diagram shows apparatus used to investigate the conductivity of different solutions.




Which substance, in aqueous solution of concentration 1 mol/dm^3 , would cause the lamp to give the brightest light?

- A ammonia
 - B ethanoic acid
 - C dilute sodium chloride solution
 - D sulfuric acid
- 27 The heat-reflecting shields of some space rockets are gold-plated, using electrolysis.
- Which electrodes and electrolyte would be used to gold-plate the heat shield?

	negative electrode	positive electrode	electrolyte
A	carbon	heat shield	gold compound
B	gold	heat shield	copper compound
C	heat shield	carbon	copper compound
D	heat shield	gold	gold compound


- 28 The following set-up shows how a block of impure copper can be purified.



The loss in mass of the anode is 50 g and the gain in mass of the cathode is 45 g.

What is the percentage purity of this sample of copper?


- A 10.0 % B 11.1 % C 90.0 % D 95.0 %
- 29 Two cells were set up as shown in the diagram. The arrow shows the direction of electron flow in the external circuit.



Which set of metals would give the electron flows in the direction shown?


	metal X	metal Y	metal Z
A	Ag	Cu	Zn
B	Ag	Zn	Cu
C	Cu	Zn	Ag
D	Zn	Cu	Ag

- 30 The energy diagram for the reaction between sodium hydroxide and hydrochloric acid is shown.



Which quantity of heat is liberated when 100 cm³ of 1 mol/dm³ hydrochloric acid reacts with 100 cm³ of 1 mol/dm³ sodium hydroxide?

- A 0.54 kJ B 2.70 kJ C 5.40 kJ D 10.8 kJ
- 31 The diagram shows the energy profile for a reaction.




Which statements about this reaction are correct?

- 1 More energy is absorbed to break the bonds than is released when new bonds are formed.
- 2 Route 1 and route 2 give the same overall equation for the reaction.
- 3 Route 2 involves the use of a catalyst.
- 4 The reaction is exothermic.

- A 1, 2 and 3 B 1 and 2 only C 2, 3 and 4 D 3 and 4 only

- 32 The equation shows a reversible reaction.



The forward reaction is endothermic.


Which of these changes will increase the yield of NO_2 ?

	pressure	temperature
A	decreased	decreased
B	decreased	increased
C	increased	decreased
D	increased	increased

- 33 Which of the following is not an example of oxidation?

- A** converting iron(III) salts to iron(II) salts
- B** converting magnesium atoms into magnesium ions
- C** dissolving a copper anode during electrolysis
- D** liberating chlorine from a chloride solution

- 34 Sulfur dioxide reacts with aqueous bromine according to the following equation.



Which element has been oxidised?

- A** bromine
- B** hydrogen
- C** oxygen
- D** sulfur


- 35 Some metals and the compounds in their ores are shown.

metal	Al	Ca	Pb	Na	Fe	Mg
compound in their ore	Al_2O_3	CaCO_3	PbS	NaCl	Fe_2O_3	MgCO_3

Which type of reaction occurs in the extraction of all of these metals from their respective ores?

- A decomposition by heat
- B electrolysis
- C precipitation
- D reduction

- 36 The diagram shows the apparatus in an experiment to reduce substance Q with the gas generated in the flask.



What are substances P and Q?

	P	Q
A	copper	copper(II) oxide
B	lead	lead(II) oxide
C	magnesium	zinc oxide
D	zinc	copper(II) oxide

- 37** Brass is an alloy.

Which statement about brass is correct?

- A** It contains a sea of electrons.
- B** It contains positive and negative ions which are free to move.
- C** It is a compound of a metal and a non-metal.
- D** It is a compound of two or more metals.

- 38** Iron is extracted from its ore haematite, Fe_2O_3 , by a reduction process in the blast furnace.

Which equation for reactions in the blast furnace shows the formation of the reducing agent?



- A** $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$
- B** $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$
- C** $\text{CO}_2 + \text{C} \rightarrow 2\text{CO}$
- D** $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$

- 39** Cars have catalytic converters fitted to reduce problems caused by some of the exhaust gases. However, cars fitted with catalytic converters still give out environmentally harmful gases.

Which of the following correctly states the harmful gas and the problem the gas causes?

	harmful gas	problem
A	nitrogen dioxide	dissolves in rain to corrode marble buildings
B	nitrogen dioxide	causes breathing problems when inhaled.
C	carbon dioxide	binds with haemoglobin in blood causing respiratory problems
D	carbon dioxide	causes the greenhouse effect leading to global warming

- 40 Nitrogen monoxide, NO, damages the ozone layer by reacting with ozone in a two-step reaction.



One nitrogen monoxide molecule can destroy thousands of ozone molecules.

Which statement correctly explains why?

- A Nitrogen monoxide in step 1 is easily generated through thunderstorms.
- B Nitrogen monoxide, while is used up in step 1, is regenerated in step 2.
- C Nitrogen dioxide produced can dissolve in rain to react with thousands of ozone molecules.
- D Nitrogen monoxide can react continuously with ozone since the ozone layer consists of thousands of ozone molecules.

End of Paper

The Periodic Table of Element


Group


I	II	Key										III	IV	V	VI	VII	0	
		1 H hydrogen 1 proton (atomic) number atomic symbol relative atomic mass																
3 Li lithium 7	4 Be beryllium 9	21 Sc scandium 45	22 Ti titanium 48	23 V vanadium 51	24 Cr chromium 52	25 Mn manganese 55	26 Fe iron 56	27 Co cobalt 59	28 Ni nickel 59	29 Cu copper 64	30 Zn zinc 65	31 Ga gallium 70	32 Ge germanium 73	33 As arsenic 75	34 Se selenium 79	35 Br bromine 80	36 Kr krypton 84	
37 Rb rubidium 85	38 Sr strontium 88	39 Y yttrium 89	40 Zr zirconium 91	41 Nb niobium 93	42 Mo molybdenum 96	43 Tc technetium -	44 Ru ruthenium 101	45 Rh rhodium 103	46 Pd palladium 106	47 Ag silver 108	48 Cd cadmium 112	49 In indium 115	50 Sn tin 119	51 Sb antimony 122	52 Te tellurium 128	53 I iodine 127	54 Xe xenon 131	
55 Cs caesium 133	56 Ba barium 137	57 – 71 lanthanoids	72 Hf hafnium 178	73 Ta tantalum 181	74 W tungsten 184	75 Re rhenium 186	76 Os osmium 190	77 Ir iridium 192	78 Pt platinum 195	79 Au gold 197	80 Hg mercury 201	81 Tl thallium 204	82 Pb lead 207	83 Bi bismuth 209	84 Po polonium -	85 At astatine -	86 Rn radon -	
87 Fr francium -	88 Ra radium -	89 – 103 actinoids	104 Rf rutherfordium -	105 Db dubnium -	106 Sg seaborgium -	107 Bh bohrium -	108 Hs hassium -	109 Mt meitnerium -	110 Ds darmstadtium -	111 Rg roentgenium -	112 Cn copernicium -	114 Fl flerovium -	116 Lv livermorium -	117 Ts tennessine -	118 Og oganesson -	119 Nh nihonium -	120 Dh dubnium -	

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm^3 at room temperature and pressure (r.t.p.).

Question	Suggested answer	Mark allocation	Remarks
1a	Ammonia	1	1e was especially badly done
b	Iodine	1	
c	Magnesium	1	
d	Argon	1	
e	Hydrochloric acid/calcium chloride	1	
2a	0.72	1	Surprisingly, a considerable number of students got this wrong
2b	A pure solvent of water is suitable for the separation of the black ink. This is because, the R_f values shows that the blue ink will travel further up the chromatogram than the red ink, While the yellow ink will remain insoluble at the original spot. Hence, all three components would be separated.	1 1	All students wrote that it is not suitable. Students need to read question carefully – since they are already told that there are only 3 dyes present in the black ink, and since two of them are soluble in pure water (with different solubilities), this means that separation is possible. Generally well done. Vague response – meet at one point
2c	When the solvent used is about 80% ethanol, all the 3 dyes will give the same R_f value as they have the same solubilities / travel the same distance. Hence, it will appear as only spot.	1 1	
2d		1m for correct set up (spot above solvent, stopper/lid) 1m for correct labelling (minimally)	Although students were not penalized, need to highlight that since ethanol, a volatile solvent, is used, a lid needs to be used to prevent it from evaporating. Need to highlight that solvent front is NOT THE SAME as start line.

			solvent, spot on pencil line)	
3a	Similarity – both are made of phosphorus atoms covalently bonded together / both involve each phosphorus atom being bonded to 3 other atoms. Difference – white phosphorus has a simple molecular structure while black phosphorus has a giant molecular / covalent structure.	1 1	Many wrote that black phosphorus has a simple molecular structure. Need to highlight that students need to name the type of bond – stating 'held together by strong bonds' is insufficient	
3b	In black phosphorus, a large amount of energy is needed to break the extensive network of strong covalent bonds. This accounts for its high melting point. However, in white phosphorus, a small amount of energy is needed to overcome the weak intermolecular forces of attraction between the molecules. Hence, it has a much lower melting point.	1 1m 1m for comparing the relative amount of energies needed.	Got ecf from 3a. Some students panicked and forgot the bonds/forces of attraction that needs to be overcome during melting.	
3c	 Key: • : electrons of Cl X : electrons of P	1m for correct bonding and arrangement of atoms	A handful of students separated the unbonded electrons in phosphorus.	


			1m for correct arrangement of electrons (-1m for drawing all electron shells)	
4a	This post is not valid. Sodium will react in air/water since it is very reactive. Hence, since rubidium is even more reactive than sodium, the possibility of it reacting in air/water is higher. Therefore, it cannot be found as silvery deposits. OR Rubidium is on top of the series. Hence it is highly reactive. It would have reacted with other elements in the air/environment to form a compound, and cannot exist uncombined as an element.	1m for explaining that it will react in air 1m for drawing comparison with sodium 1 1	Generally well done	
4b	Magnesium reacts very slowly with cold water while iron does not react with cold water. Since beryllium is in between these two metals, it might not even react / react extremely slowly (slower than Mg) with cold water Magnesium reacts violently with steam whereas iron reacts slowly with steam. Since beryllium is between these two metals, it might react quickly/rapidly/slowly (slower than Mg) with steam.	1 1 1 1	Poorly done. Need to highlight that the question's intent is for them to explain the vigour of the reaction, and also that they need to draw comparisons with magnesium and iron, since these two are the located the closes with beryllium.	
4c	Heating bismuth oxide with coke (carbon) / hydrogen			Some wrote 'electrolysis'
5a(i)	Ba^{2+} , OH^-	1		To go through 5a and 5b in detail since it was very badly done.
5a(ii)	H^+ , SO_4^{2-}	1		Even for 5a, many students stated the formula of the substance though they have been clearly
5b	At the start of the experiment, the ammeter reading was the highest as there was highest number of ions present (concentration of ions is at its maximum) to carry a current.	1		

	As sulfuric acid is added, the ammeter reading drops as the Ba^{2+} ions were used up to form solid BaSO_4 . Therefore, there are fewer mobile ions present to carry a current. When 5.00 cm^3 of sulfuric acid is added, all the Ba^{2+} ions have reacted to form solid barium sulfate, which is poor electrical conductor.	1	asked for formula of the ions. Many failed to interpret the curve correctly.
5c	No. of moles of sulfuric acid needed for complete reaction = $0.500 \times 0.005 = 0.0025$ Concentration = $0.0025 / 0.025 = 0.1 \text{ mol/dm}^3$ OR $M_1V_1 / M_2V_2 = n_1/n_2$ $M_1 \times 25 / (0.5 \times 5) = 1/1$ $M_1 = 0.1 \text{ mol/dm}^3$	1 1 1 1 1	Generally well done.
5d	Plastic prevents the iron from coming into contact with oxygen and water, preventing rusting.	1	Generally well done. Vague response will be to just state that iron will react with the mixture.
6a	Zn^{2+} or Cu^{2+} (zinc or copper(II) ions) As aqueous ammonia is added, a precipitate of the metal hydroxide forms, and mass of solid increases. When aqueous ammonia is added in excess, the precipitate dissolves and the mass of solid decreases.	1 1 1	Very poorly done. Need to highlight to students that since question is asking them to identify cation, they need to zoom in straight to the cations test.
6b	No. The reaction of the metal oxide obtained from stage 1 with hydrochloric acid will not produce carbon dioxide.	1	Many failed to give the proper explanation, just stating that there is 'no more carbonate left.'

6c	$\text{Ag}^+(\text{aq}) + \text{Cl}^-(\text{aq}) \rightarrow \text{AgCl}(\text{s})$		1m for correct balanced equation, 1m for correct state symbols	Poorly done. Tried to award ecf but tough..								
7a	This is to ensure that all of the oxygen has reacted.		1	Generally well done. Vague responses will include 'ensuring that all reactants has reacted'.								
b	Volume of oxygen = $200 - 160 = 40 \text{ cm}^3$ % of oxygen = $40 / 200 \times 100\% = 20\%$		1									
c(i)	<table><tr><td></td><td>oxidation state of carbon</td></tr><tr><td>FeC_2O_4</td><td>+3</td></tr><tr><td>CO_2</td><td>+4</td></tr><tr><td>CO</td><td>+2</td></tr></table>		oxidation state of carbon	FeC_2O_4	+3	CO_2	+4	CO	+2		1m for FeC_2O_4 1m for the other 2 substances	Very poorly done for FeC_2O_4
	oxidation state of carbon											
FeC_2O_4	+3											
CO_2	+4											
CO	+2											
c(ii)	The oxidation state of C has increased from +3 (in FeC_2O_4) to +4 (in CO_2). The oxidation state of C has also decreased from +3 (in FeC_2O_4) to +2 (in CO). Since both oxidation and reduction happens at the same time, this is a redox reaction.		1 1	Decent								
8a	30%		1	Well done								

8b	This is to increase the yield of hydrogen and nitrogen into ammonia. This also helps to save cost and/or prevent wastage.			1	Well done
8c	When the temperature is too low, the speed of reaction will be too slow.			1	Well done
9a		cathode	anode	1 each	Extremely poorly done. Really need to teach students to read and process passage
	AFC	$O_2(g) + 2H_2O(l) + 4e \rightarrow 4OH^-(aq)$	$H_2(g) + 2OH^-(aq) \rightarrow 2H_2O(l) + 2e$		
	PEMFC	$4H^+(aq) + O_2(g) + 4e \rightarrow 2H_2O(l)$	$H_2(g) \rightarrow 2H^+(aq) + 2e$		
9b	$2H_2(g) + O_2(g) \rightarrow 2H_2O(l)$			1 for correct balanced equation, 1 for state symbols	Poorly done
9c	Carbon dioxide, being an acidic oxide, reacts with the electrolyte potassium hydroxide.			1	Even though many were able to state that carbon dioxide is an acidic oxide, many failed to write the correct equation; instead they wrote the equation for the formation of carbonic acid, which doesn't explain why carbon dioxide poisons the electrolyte at all.
	$CO_2 + 2KOH \rightarrow K_2CO_3 + H_2O$			1	
9d	The operation of AFC requires non-precious metal as catalyst in contrast to PEMFC which requires expensive platinum catalysts and precious metal-based electrodes as well.			1	Well done.

9e(i)	They produce only water as the waste product .	1	Many wrote 'they do not produce greenhouse gases'; there are many other ways that a fuel can pollute the environment.
9e(ii)	Methane is used in the production of hydrogen. If methane leaks into the atmosphere, it can trap heat and cause greenhouse effect . OR Carbon monoxide, which is produced, is toxic as it binds to the haemoglobin, resulting in it being unable to bind to oxygen, which may lead to breathing difficulties that may result in death.	1	Students need to discuss the impact. For CO, many just stated that it is a toxic gas without explaining its impact.
10a	$2\text{H}_2\text{O}_2 \rightarrow \text{O}_2 + 2\text{H}_2\text{O}$	1	Many did not know the formula of hydrogen peroxide!
10b	More energy is released to form O=O bonds and O-H bonds than the energy taken in to break O-H bonds and O-O bonds. OR Less energy is taken in to break O-H bonds and O-O bonds than the energy released to form O=O bonds and O-H bonds.	1 1 1	Many did not split up the atoms. For e.g 'H-O-H bonds'.

10c(i)		1m for correct shape of energy profile diagram, 1m for labelling the activation energy on both energy profile diagrams, 1m for correctly labelling the reactants and products	Generally well done.
10c(ii)	The catalyst lowers the activation energy of the reaction and this increases the number of particles having energy higher than or equal to the new activation energy. Hence, the frequency of effective collisions increases, thereby speeding up the reaction.	1 1	Many failed to realise that since this is a part question, their answer needs to be linked to c(i).
10c(iii)	increase the concentration of the hydrogen peroxide	1	Many wrote 'increase temperature' without considering the feasibility of it in this context.
EITHER 11a(i)	$2\text{NiS} + 3\text{O}_2 \rightarrow 2\text{NiO} + 2\text{SO}_2$	1	A few did not balance correctly.
11a(ii)	No. of moles of NiS = $182\,000 / 91 = 2000$ No. of moles of SO_2 formed = 2000 Volume = $2000 \times 24\text{ dm}^3 = 48\,000\text{ dm}^3$	1 1	Many did not convert the mass to grams! Ecf awarded (grudgingly)

11b	Metals consist of a lattice of positive ions that are surrounded by a sea of mobile delocalized free moving electrons . This sea of mobile electrons can move to carry the charge of an electric current when a potential difference is applied.	1 1	Well done. Those who did not get the full marks did not explain metal structure.
11c	No visible reaction Blue solution turns green Nickel gets coated with a pink solid	1 1 1	Many wrote 'blue solution turns colourless'
11d	Magnesium is more reactive than nickel /carbon Its oxide cannot be reduced by carbon to form magnesium ; however, nickel(II) oxide can be reduced by carbon in stage 2. OR Magnesium can only be obtained by electrolysis	1 1	Need to highlight to students that they need to compare magnesium to either Ni or C.
OR 11a	$\text{NaOH(aq)} + \text{H}_3\text{PO}_4\text{(aq)} \rightarrow \text{NaH}_2\text{PO}_4\text{(aq)} + \text{H}_2\text{O(l)}$ Magnesium can only be obtained by electrolysis	1m for correct equation, 1m for correct state symbols	Not many attempted this question.
11b(i)	Strong acid – sulfuric acid Weak acid – phosphoric acid / sodium dihydrogen phosphate / ethanoic acid Sulfuric acid is a strong acid as it undergoes complete ionisation in water that results in a high concentration of hydrogen ions , thereby resulting in a low pH. Phosphoric acid / sodium dihydrogen phosphate / ethanoic acid is a weak acid as it undergoes partial ionisation in water to form a low concentration of hydrogen ions , thereby resulting in a relatively high pH.	1m for correctly stating the types of ionisation that strong and weak acids undergo 1m for stating that ionization takes place in water 1m for stating the relative concentrations of hydrogen ions formed	Many missed out 'in water'

